

The Silent Biliary Dysfunction Hypothesis

ABSURD HEALTH INSTITUTE · 03 · Hepatobiliary, cholesterol and endocrine physiology · 15 OCTOBER 2026

Abstract

A patient may report food-related discomfort, altered bowel habits or intolerance to fatty meals without an obvious gallstone or duct obstruction. Could selected functional disturbances of biliary physiology contribute to these complaints before structural imaging becomes abnormal? This paper defines a narrower, testable concept of biliary dysfunction and rejects the idea that an invisible blockage should be presumed whenever symptoms remain unexplained.

The Clinical Question and Its Context

Biliary physiology spans hepatocyte bile-acid synthesis, conjugation, canalicular secretion, duct transit, gallbladder storage and contraction, pancreatic coordination, intestinal micelle formation and ileal reuptake. A defect at one step can cause a different syndrome from failure at another. Cholestasis is not synonymous with gallbladder dyskinesia; bile-acid diarrhea is not the same as insufficient luminal bile acids; functional abdominal pain may occur without impaired bile delivery. Nonspecific pale or variable stools cannot reliably quantify enterohepatic function.

Proposed Biological or Explanatory Model

The construct becomes useful if phenotypes are defined: clinically significant cholestasis with biochemical markers; documented gallbladder emptying abnormalities in an appropriate setting; bile-acid malabsorption; pancreatic insufficiency; or established intestinal fat malabsorption. The analysis should use symptom chronology, medication history and targeted testing rather than the unvalidated term “stagnant bile” as a diagnosis. Functional disturbances may be real, but proving them requires more than the absence of structural findings.

Mechanism-Specific Interpretation

The phrase “silent dysfunction” is most useful if it describes a patient group with a reproducible physiologic abnormality, not a residual category for symptoms medicine has not yet explained. A person with postprandial abdominal pain and objectively impaired gallbladder emptying is not in the same group as a patient with normal imaging, normal blood tests and diffuse fatigue. Likewise, bile-acid diarrhea entails excess bile acids reaching the colon rather than necessarily deficient bile in the small intestine. Conflating these states would produce mutually contradictory treatment assumptions. Precision begins by describing the symptom, organ, proposed mechanism and evidence that the mechanism is present.

Competing Explanations and Limits of Inference

Fatty-meal discomfort also occurs with reflux, dyspepsia, peptic disease, functional gastrointestinal disorders and food intolerance. Gallbladder imaging is imperfect, but negative imaging does not establish a hidden sludge process. HIDA scan ejection fractions have interpretive limits and should not become a universal screen. Supplement responses are vulnerable to placebo effects and regression to the mean.

Discriminating Observations and Evidence Thresholds

The decisive question is whether functional measurements explain outcomes after standard gastrointestinal diagnoses and medication effects are considered. If the proposed biomarkers do not distinguish symptomatic from asymptomatic people, “silent biliary dysfunction” has not yet gained clinical meaning. A successful concept must yield a defined phenotype and useful discriminating information, not merely an evocative name.

Implications for Interpretation and Future Inquiry

The proposed clinical outcome is a better diagnostic taxonomy, not universal prescription of bitters, bile salts or “liver flushes.” Cholestatic jaundice, severe right-upper-quadrant pain, fever or dark urine with pale stools require conventional medical assessment. A carefully defined functional phenotype could invite future physiologic study without replacing established gastroenterology.

Selected Evidence and Further Reading

1. [Recent Advances in the Digestive, Metabolic and Therapeutic Effects of FXR and FGF19](#)
2. [Bile acids: regulation of synthesis](#)

Source lineage: Terrain Medicine Handbook; Bile and hormonal investigations

Editorial note: Conceptual scientific commentary; not an original clinical trial, systematic review, peer-reviewed publication, or medical advice. Hypotheses are not established treatment effects.