

# Chronic Hyponatremia and the Misdiagnosis of Neurological Decline

ABSURD HEALTH INSTITUTE | 15 OCTOBER 2026 | IV. The diluted brain and osmotic neurology

---

## Abstract

This concept paper examines chronic hyponatremia and the misdiagnosis of neurological decline. It develops an argument drawn from *The Diluted Mind*, identifies the relevant biological or clinical mechanisms, and considers where that argument may fail. This is a narrative perspective, not a systematic review or an original clinical study.

## The central problem

Chronic mild-to-moderate hyponatremia has often been described as relatively asymptomatic because dramatic edema is less common after osmotic adaptation. That does not make it functionally trivial. The research question is whether sustained low sodium can contribute to impaired gait, attention and falls that might otherwise be attributed exclusively to aging, medication, depression or neurodegeneration.

## Mechanisms and causal architecture

Renneboog and colleagues examined associations between chronic hyponatremia and falls and reported changes in gait and attention testing that improved after correction in a small subset. This is a clinically suggestive finding, not proof that all subtle neurological decline is hyponatremic. Older adults have multiple overlapping causes of imbalance: neuropathy, visual impairment, vestibular disease, sedating medications, frailty and cerebrovascular disease. The plausible neurological mechanisms include altered cellular osmolytes, neurotransmission and network performance, but direct causal attribution in a particular individual requires temporal and clinical correlation.

## A discriminating example

A patient in their seventies develops worsening attention and falls while taking a thiazide diuretic. Chronic hyponatremia may contribute, but neuropathy, orthostasis, vision and medications also warrant evaluation. The paper can show why correcting an electrolyte abnormality under clinical supervision may improve some functions without claiming that every apparent decline was caused by dilution.

## Competing explanations and limitations

Hyponatremia can be a marker of more severe illness, diuretic exposure, malignancy, endocrine dysfunction or heart failure. Observational associations may reflect these confounders. Some patients with chronic sodium abnormalities have no measurable cognitive impairment. Correcting sodium is not automatically a treatment for dementia, and correction itself carries risk in susceptible chronically adapted patients.

## What evidence would change the interpretation?

The relevant observations are onset and duration of low sodium, medications, paired urine studies, neurological examination and changes in validated gait or attention outcomes as sodium normalizes under appropriate supervision. A genuine association should be reproducible after adjustment for confounders, with clear clinical effect sizes rather than merely significant laboratory correlations.

## Implications for medical interpretation

The clinical lesson is to include sodium and tonicity in the differential for otherwise unexplained gait or cognitive decline, while continuing to evaluate neurological and geriatric causes. No rapid home correction or empirical salt loading follows from this argument.

## Selected scientific reading

Renneboog et al. (2006), Mild chronic hyponatremia and falls, attention deficits, American Journal of Medicine.  
<https://doi.org/10.1016/j.amjmed.2005.09.026>

Spasovski et al. (2014), Clinical practice guideline on diagnosis and treatment of hyponatraemia, European Journal of Endocrinology.  
<https://doi.org/10.1530/EJE-13-1020>

Adaptation of the Brain to Hyponatremia and Its Clinical Implications (2023), Journal of Clinical Medicine.  
<https://doi.org/10.3390/jcm12051714>

Original source: The Diluted Mind. All source books supplied by the author.

Conceptual research commentary. Not a peer-reviewed clinical study, original trial, systematic review, validated diagnostic instrument or treatment protocol. Consult qualified care for medical decisions.